

**ANALISIS DESKRIPTIF TERHADAP KASUS DATA PERSALINAN DI
BANGSAL OBGYN PADA TRIWULAN IV RUMAH SAKIT PANTI
WILASA Dr.CIPTO SEMARANG PERIODE 2011**

DEDY ARISANDI

*Program Studi Rekam Medis&Info. Kesehatan - D3, Fakultas
Kesehatan, Universitas Dian Nuswantoro Semarang*

URL : <http://dinus.ac.id/>

Email : derianonk@yahoo.co.id

ABSTRAK

Sampai sekarang masalah kesehatan ibu dan anak masih merupakan masalah pokok yang dihadapi bangsa Indonesia termasuk kasus persalinan yang semakin bervariasi. Data menunjukkan angka bedah caesarea di rumah sakit swasta diatas 30% bahkan ada yang mencapai 80%. Di RS Panti Wilasa dr. Cipto Semarang pada triwulan 4 tahun 2011 mencapai 74%. Penelitian bertujuan untuk mengetahui deskripsi jenis persalinan pada pasien yang menjalani persalinan di bangsal obsgyn RS Panti Wilasa dr. Cipto Semarang.

Penelitian yang digunakan adalah penelitian deskriptif dengan pendekatan retrospektif . Populasi dari penelitian ini adalah data rekam medis pasien yang menjalani persalinan di bangsal obsgyn RS Panti Wilasa dr. Cipto Semarang pada triwulan 4 tahun 2011. Sampel penelitian adalah seluruh populasi yaitu 123 pasien (total sampling). Data yang digunakan berupa data sekunder yang diperoleh dari observasi terhadap data RM 1 pasien. Analisa data dilakukan secara deskriptif dalam bentuk table, grafik dan narasi.

Dari 123 kasus persalinan pada tahun 2011, terdapat 25% kasus persalinan normal, 1% kasus persalinan vakum ekstraksi dan 74% kasus persalinan sectio caesarea. Umur ibu yang melakukan persalinan yaitu umur <20 tahun 24%, umur 20-35 tahun 63% dan umur >35 tahun 13%. Diagnosa utama dalam sebuah persalinan sangat menentukan dalam penatalaksanaan akhir dari sebuah persalinan, diagnose utama yang ada pada penelitian ini terdiri dari bekas sectio caesarea (14%), Ketuban Pecah Dini (13%), Partus Tak Maju (16%), Pre Eklamsi Berat (10%), Induksi gagal (8%), Fetal Distress (10%). Terdapat pula diagnosa lain ditentukan untuk mendukung diagnosa utama Pada penelitian ini diagnosa lain terdiri dari kasus Cephalopelvic Disproportion (CPD) (15%), Anemia (5%), Diabetes Mellitus (2%), Ketuban Pecah Dini (35%), Partus Tak Maju (23%), Pre Eklamsi Berat (13%).

Dari hasil penelitian disarankan perlu meneliti lebih lanjut melalui pembuktian secara statistik tentang factor-faktor yang berhubungan dengan jenis persalinan serta adanya kerja sama antara tenaga kesehatan dengan pasien dalam pemberi pelayanan.

Kata Kunci : Analisis Deskriptif, Jenis Persalinan, Usia Ibu, Diagnosa Utama Dan Diagnosa Lain

**DESCRIPTIVE ANALITIC TO THE 4-TH QUARTERLY DATA OF
DELIVERY CASES AT OBSTETRY GYNECOLOGY DEPARTMENT IN
PANTI WILASA dr. CIPTO
SEMARANG HOSPITALS 2012**

DEDY ARISANDI

*Program Studi Rekam Medis&Info. Kesehatan - D3, Fakultas
Kesehatan, Universitas Dian Nuswantoro Semarang*

URL : <http://dinus.ac.id/>

Email : derianonk@yahoo.co.id

ABSTRACT

Hitherto, the health issue of mother and child is the main issue that Indonesia deal with, including delivery cases which are increasingly varied. It is characterized by the high rate of SC delivery in Panti Wilaya Dr. Cipto Semarang hospital.

This research was conducted by using descriptive study and retrospective approach. The population of this study is medical record of 123 maternals who underwent the childbirth process in Obstetry Gynecology department of Panti Wilasa Dr. Cipto Semarang Hospital, especially 4th quarterly in 2011 which engaged the stated criteria i.e 1st form of medical record or the data of the client completely recorded. Sample for this research is all the population by using total sampling method. The data that used in this study is secunder data that obtained from the directly observation to the medical record. The data was tabulated by computerization and by using univariat data analysis.

Among 123 cases of delivery in 2011, there are 31 cases of normal delivery, 1 case of vacuum extraction delivery and 91 cases of sectio caesarea delivery. Sectio Caesarea, among delivery methods, is the most high case of delivery method. The maternal age in this study were grouped to 3 categories i.e 29 are mothers in 15 - 25 years old group, 78 are mothers in 26 - 35 years old group and 16 mothers are in 36 - 45 years old group. Maternal age is one of predisposition factors which influence delivery proses of a mother. Management of labor depends on the main diagnosis of a delivery. Some of the main diagnoses in this study were as follows 17 cases of previously SC, 16 cases of premature rupture membrane, 20 cases of secondary arrested/prolonged secondary stage, 12 cases of pre-eclampsia, 10 cases of failed induction, 12 cases of feotal distress and 36 cases of others i.e normal and pathology delivery. There are another diagnoses which support the primary diagnosis in this study. These diagnoses are 18 cases of cephalopelvic disproportion, 7 cases of postpartum anemia, and 2 cases of gestational DM.

The conclusion Of this reaserch generally is that the description to the 4th quarterly of delivery cases in Obsgyn departement of Panti Wilasa Dr. Cipto Semarang Hospital mostly implemented by SC. This is supported by the factor of maternal age, main diagnosis, supporting diagnosis and the prosedure of delivery which is chosen by the qualified medical and non-medical team.

Keyword : Analisis Deskriptif, Jenis Persalinan, Usia Ibu, Diagnosa Utama Dan Diagnosa Lain